



A tumor-targeted CD40 agonistic DARPin® molecule leading to antitumor activity with limited systemic toxicity

Nicolò Rigamonti, Ph.D.

Molecular Partners AG, Schlieren-Zurich, Switzerland

Disclosure Information

I have the following financial relationships to disclose:


Employee of: Molecular Partners AG, Schlieren-Zurich, Switzerland

- and -


I will not discuss off label use and/or investigational use in my presentation.

DARPin® platform

Targeting peptide
MHC complexes
Poster 690




Tumor-localized immune
cell modulation
Poster 2273



Next-generation
T-cell engagers

DARPin® Features

Rigid-body target binding




DARPin® domain

Multi-DARPin® formatting



SCALE

15 kDa



150 kDa

- High specificity and affinity
- Small size: 15 kDa
- Simple repetitive architecture: 1 polypeptide
- Tunable half-life

MP0317, a multi-specific DARPin® molecule, designed to improve efficacy and safety via tumor localized CD40 agonism




FAP
(fibroblast activation protein)




tumor localization and clustering

CD40




immune cell modulation

HSA
(human serum albumin)




systemic half-life extension




FAP expression in human tumor sections


Colon




Lung




Stomach




Uterus



Ovary




Pancreas




Human FAP, DAPI

MP0317, a FAP-dependent MoA across different immune cell populations



Dendritic cells:

CD86 ✓ IL-12 ✓
CD83 ✓ CD80 ✓






Macrophages:

TNF α ✓ IL-12 ✓
IL-6 ✓ CCL17 ✓



B cells:


CD69 ✓
CD86 ✓




Mouse-specific FAP x CD40 DARPin® molecule (mFAP x mCD40) localizes and accumulates to the FAP- transfected MC38 tumor (MC38-FAP)



A
SPECT-CT study
(mFAP x mCD40)




B
Detection of the DARPin®
in the tumor by IHC




- CTRL DARPin® binding HSA, but neither FAP nor CD40

C
Biodistribution study






mFAP x mCD40 restrains tumor progression in a FAP-dependent manner




mFAP x mCD40 provides durable anti-tumor response in animals bearing FAP-transfected MC38 tumors




MC38-FAP
Colorectal cancer



mFAP x mCD40 induces a protective anti-tumor immunological memory that is not limited to the FAP antigen




In the TME, mFAP x mCD40 activates DC, increases the presence of M1-like macrophages and lymphocytes



Differently than CD40 Ab, mFAP x mCD40 does not show signs of systemic toxicity

A

Lack of weight lost




Time after tumor inoculation (days)

Vehicle
Neg. CTRL
mFAP x mCD40
mCD40 Ab







B

Lack of increase of proinflammatory cytokines in the serum



C

Lack of transaminase elevation in the serum and hepatocellular changes such as immune cell infiltration (ICI) and necrosis (NEC).



Conclusions

- Double targeting FAP x CD40 DARPin® molecules activate in a FAP dependent manner different immune cell populations *in vitro* and induce a strong and durable antitumor response in FAP-positive tumors *in vivo* without showing signs of systemic toxicity
 - MP0317 activates *in vitro* 1) monocyte-derived dendritic cells 2) monocyte-derived macrophages and 3) B cells in a strictly FAP-dependent manner
 - Mouse mFAP x mCD40 shows a strong antitumor activity in FAP^{HIGH} tumor-bearing mice, but has a negligible antitumor effect on FAP^{NEG} or FAP^{LOW} neoplasias confirming a FAP-specific MoA *in vivo*
 - mFAP x mCD40 generates anti-tumor immune memory response
 - mFAP x mCD40 affects the DC and macrophages and increases the percentage of lymphocytes in the TME
 - mFAP x mCD40, differently than the CD40 Ab, does not show signs of systemic toxicity

Acknowledgement



MOLECULAR
partners

Andrea Valeri,
Anja Schlegel,
Clara Metz,
Christof Zitt,
Heidi Poulet,
Jonas Schwestermann,
Niels Breitenstein,
Niina Veitonmäki,
Omar Abdelmotaleb,
Ralph Bassey,

Pamela A. Trail,
Sophie Barsin,
Susanne Mangold,
Yvonne Kaufmann,
Victor Levitsky,
And all colleagues at
Molecular Partners

PAUL SCHERRER INSTITUT



Alain Blanc,
Martin Behé,
Roger Schibli,
Stefan Imobersteg